

AN ANALOGUE OF SELBERG'S FORMULA FOR MOTOHASHI'S PRODUCT

Sergei N. Preobrazhenskii¹

We prove an analogue of Selberg's explicit formula for Motohashi's product (see [arXiv:1104.1358v3 \[math.NT\]](#)). We also provide a zero-density theorem for the product, which follows from Soundararajan's theorem for moments of the Riemann zeta-function on the critical line.

Key words: Riemann zeta-function, Riemann hypothesis.

1. Introduction. In [1] we gave a proof of the following theorem.

Assume RH. Then there exists an infinite sequence of pairs of real numbers (T_1, T_2) , $T_1 = T$, $T_2 = T + H$, with arbitrarily large values of T and $H = c(\log \log T)^{-1}$, such that

$$|\zeta(1 + iT_1)| |\zeta(1 + iT_2)| \ll (\log \log T)^{-2}$$

and

$$\begin{aligned} & (\log \log T)^7 |\zeta(1 + iT_1)|^4 |\zeta(1 + iT_2)|^4 \\ & + (\log \log T)^7 \zeta(1 + i(T_1 + H))^2 \zeta(1 - i(T_1 - H))^2 \zeta(1 + i(T_2 + H))^2 \zeta(1 - i(T_2 - H))^2 \\ & + (\log \log T)^7 \zeta(1 + i(T_1 - H))^2 \zeta(1 - i(T_1 + H))^2 \zeta(1 + i(T_2 - H))^2 \zeta(1 - i(T_2 + H))^2 \\ & \ll (\log \log T)^{-1}. \end{aligned}$$

We put

$$z = \exp(A \log \log T \log \log \log T)$$

with the same constant A as in [1], set

$$\xi(d) = \lambda_d(z) = \begin{cases} \mu(d) & \text{if } d < z, \\ \mu(d) \frac{\log(z^2/d)}{\log z} & \text{if } z \leq d < z^2, \\ 0 & \text{otherwise,} \end{cases}$$

$$P_d(s, T_1, T_2)$$

$$= \prod_{p|d} \left(1 - \left(1 - \frac{1}{p^s} \right) \left(1 - \frac{1}{p^{s-iT_1}} \right) \left(1 - \frac{1}{p^{s+iT_2}} \right) \left(1 - \frac{1}{p^{s-i(T_1-T_2)}} \right) \left(1 - \frac{1}{p^{2s-i(T_1-T_2)}} \right)^{-1} \right),$$

and write

$$\begin{aligned} J(s, T_1, T_2) &= \frac{\zeta(s) \zeta(s - iT_1) \zeta(s + iT_2) \zeta(s - i(T_1 - T_2))}{\zeta(2s - i(T_1 - T_2))}, \\ K(s, T_1, T_2) &= \sum_{d \leq z^2} \lambda_d(z) P_d(s, T_1, T_2). \end{aligned}$$

The theorem below shows that on RH the absolute value of the product JK cannot exceed a quite small function $((\log \log T)^\epsilon)$ in a quite large neighborhood (within the distance $c(\log \log T)^{-1}$) to the left of the line $\Re s = 1$.

¹*Preobrazhenskii Sergei Nikolayevich* — Department of Mathematical Analysis, Faculty of Mechanics and Mathematics, Lomonosov Moscow State University.

Theorem 1. Assume the Riemann hypothesis. Let $s_0 = \sigma_0 + it_0$ be a point such that

$$|J(s_0, T_1, T_2)K(s_0, T_1, T_2)| \geq (\log \log T)^\varepsilon$$

with arbitrarily small fixed $\varepsilon > 0$, and

$$\sigma_0 = 1 - \frac{E_0}{\log \log T} \geq 1 - \frac{E}{\log \log T}, \quad C \log \log \log T \leq |t_0| \leq T/2. \quad (1)$$

Then $E_0 \geq c_2(\varepsilon) > 0$.

Here we call the product $J(s_0, T_1, T_2)K(s_0, T_1, T_2)$ Motohashi's product (see [2]).

In this paper, we prove an analogue of Selberg's formula for the logarithmic derivative of Motohashi's product. We also provide a zero-density theorem for the product, which follows from Soundararajan's theorem for moments of the Riemann zeta-function on the critical line [3].

2. Lemmas and the results. The following lemmas are found in Ingham [4], theorems 26 and 27. The proofs use the functional equation for the zeta-function.

Lemma 1. There exists a sequence of numbers $T_2, T_3, \dots$, such that

$$m < T_m < m + 1 \quad (m = 2, 3, \dots)$$

and

$$\left| \frac{\zeta'}{\zeta}(s) \right| < A \log^2 t \quad (-1 \leq \sigma \leq 2, t = T_m).$$

Lemma 2. In the region obtained by removing from the half-plane $\sigma \leq -1$ the interiors of a set of circles of radius $\frac{1}{2}$ with centres at $s = -2, -4, -6, \dots$, i.e. in the region defined by

$$\sigma \leq -1, \quad |s - n| \geq \frac{1}{2} \quad (n = -2, -4, -6, \dots),$$

we have

$$\left| \frac{\zeta'}{\zeta}(s) \right| < A \log(|s| + 1).$$

The following lemma is a consequence of lemmas 1 and 2 (cf. Lemma 9 of [5]).

Lemma 3. There exists a sequence of numbers $T_2, T_3, \dots$, such that

$$m < T_m < m + 1 \quad (m = 2, 3, \dots)$$

and

$$\left| \frac{\zeta'}{\zeta}(s) \right| < A \log^2 m,$$

for $\sigma \geq -m - 1/2$, $t = \pm T_m$, or $\sigma = -m - 1/2$, $|t| < T_m$.

Consider the function $Z(s) = J(s, T_1, T_2)K(s, T_1, T_2)$. By Lemma 3 of [1], for $\sigma > 1$ $Z(s)$ is given by the convergent Dirichlet series

$$Z(s) = \sum_{n=1}^{\infty} \sigma_{iT_1}(n) \sigma_{-iT_2}(n) \left(\sum_{d|n} \xi(d) \right) n^{-s}.$$

Recalling the definition of $\xi(d)$ we see that for σ sufficiently large we can make the power series expansion for $\log Z(s)$ as $\log(1 + x)$ and represent this as the convergent Dirichlet series.

Alternatively, we can expand $\log K(s)$ and use the known expansion for $\log J(s)$. Differentiating term by term, we obtain the Dirichlet series for $\frac{Z'}{Z}(s)$. Denote the coefficients of the series by $\Sigma(n)$:

$$\frac{Z'}{Z}(s) = \sum_{n=2}^{\infty} \frac{\Sigma(n)}{n^s}.$$

The following formula is an analogue of Selberg's formula for $\frac{\zeta'}{\zeta}(s)$. We denote the complex zeros of $\zeta(s)$ by $\rho = \beta + i\gamma$, the zeros of $K(s, T_1, T_2)$ by r , and the poles of $K(s, T_1, T_2)$ by ν .

Theorem 2. *Let $x > 1$ and write*

$$\Sigma_x(n) = \begin{cases} \Sigma(n) & \text{for } 1 \leq n \leq x, \\ \Sigma(n) \frac{\log^2 \frac{x^3}{n} - 2 \log^2 \frac{x^2}{n}}{2 \log^2 x} & \text{for } x \leq n \leq x^2, \\ \Sigma(n) \frac{\log^2 \frac{x^3}{n}}{2 \log^2 x} & \text{for } x^2 \leq n \leq x^3. \end{cases}$$

Then for

$$s \notin S_1 = \{1, 1 + iT_1, 1 - iT_2, 1 + i(T_1 - T_2)\}, \quad s \notin \overline{S}_1 = \left\{ \frac{1}{2} + \frac{i}{2}(T_1 - T_2) \right\},$$

$$s \notin S_\rho = \{\rho, \rho + iT_1, \rho - iT_2, \rho + i(T_1 - T_2)\}, \quad s \notin \overline{S}_\rho = \left\{ \frac{\rho}{2} + \frac{i}{2}(T_1 - T_2) \right\},$$

$$s \notin S_{-2q} = \{-2q, -2q + iT_1, -2q - iT_2, -2q + i(T_1 - T_2)\}, \quad s \notin \overline{S}_{-2q} = \left\{ -q + \frac{i}{2}(T_1 - T_2) \right\}$$

$$(q = 1, 2, 3, \dots),$$

$$s \notin S_\nu = \left\{ \nu = i \left(\frac{\pi k}{\log p} + \frac{1}{2}(T_1 - T_2) \right) \quad (k \in \mathbb{Z}' \subset \mathbb{Z}, \quad p < z^2) \right\},$$

$$s \notin S_r = \{r\},$$

we have

$$\begin{aligned} \frac{Z'}{Z}(s) = & \sum_{n < x^3} \frac{\Sigma_x(n)}{n^s} + \sum_{u \in S_1 \cup \overline{S}_\rho \cup \overline{S}_{-2q} \cup S_\nu} \frac{x^{u-s}(1 - x^{u-s})^2}{\log^2 x(u-s)^3} \\ & - \sum_{u \in \overline{S}_1 \cup S_\rho \cup S_{-2q} \cup S_r} \frac{x^{u-s}(1 - x^{u-s})^2}{\log^2 x(u-s)^3}, \end{aligned} \quad (2)$$

where the zeros and the poles are counted according to their multiplicities.

For the proof, we follow the argument of Selberg in Lemma 10 of [5]. Note that the analogue of Lemma 3 for the function $K(s, T_1, T_2)$ should be with the bound

$$\left| \frac{K'}{K}(s, T_1, T_2, z) \right| < A_{T_1, T_2, z} m, \quad (3)$$

for a sequence of contours C_m : $\sigma_m \leq \sigma \leq \alpha$, $t = \pm T_m$, or $\sigma = \sigma_m$, $|t| < T_m$ ($\sigma_m \rightarrow -\infty$, $T_m \rightarrow +\infty$). The sequence σ_m, T_m is chosen properly so as to avoid zeros and poles of K (and zeros of J). This can be done using Jensen's theorem for K . In view of the bound (3) we need degree 3 in the denominator for the integral to tend to zero.

Next, we give a zero-density theorem for Motohashi's product. Recall that

$$K_X(s) = K(s, T_1, T_2) = \sum_{d \leq X^2} \lambda_d(X) P_d(s, T_1, T_2),$$

and

$$J(s)K_X(s) = 1 + \sum_{n=X}^{\infty} \sigma_{iT_1}(n) \sigma_{-iT_2}(n) \left(\sum_{d|n} \lambda_d(X) \right) n^{-s}.$$

Let

$$f_X(s) = J(s)K_X(s) - 1,$$

and define $N_K(\sigma, T)$ to be the number of zeros r of the function $K_X(s)$ such that $\Re r > \sigma \geq \frac{1}{2}$, $0 < \Im r \leq T$.

Lemma 4. *If for some $X = X(\sigma, T)$, $X < T^{A_1}$,*

$$\int_0^T |f_X(\sigma + it)|^{1/\varkappa} dt = O(T^{l(\sigma)} \log^m T)$$

as $T \rightarrow \infty$, uniformly for $\alpha \leq \sigma \leq \beta$, where $l(\sigma)$ is a positive non-increasing function with a bounded derivative, $\varkappa$ lies between $\frac{1}{2}$ and $\frac{3}{4}$, and m is a constant ≥ 0 , then for any fixed $A_2 \geq 1$

$$N_K(\sigma, T) = O(T^{l(\sigma)} \log^{m+1} T) + O\left(\frac{T \log T}{X^{A_2}}\right)$$

uniformly for $\alpha + 1/\log T \leq \sigma \leq \beta$.

This lemma is a modified form of Theorem 9.16 of [6].

Lemma 5. *Assume RH. For every positive real number k and every $\varepsilon > 0$ we have*

$$\int_0^T \left| \zeta\left(\frac{1}{2} + it\right) \right|^{2k} dt \ll_{k,\varepsilon} T(\log T)^{k^2+\varepsilon}.$$

This is a theorem of K. Soundararajan [3].

The following lemma is an analogue of Theorem 9.19 (B) of [6] (Ingham's zero density theorem) and follows from Soundararajan's theorem and the upper bound for $K_X(s)$ with $X = z$ in [1].

Lemma 6. *Assume RH.*

$$\int_0^T |f_X(1/2 + d \log \log T \log \log \log T / \log T + it)|^{1/\varkappa} dt = O(T^{1-c_d \log \log T \log \log \log T / \log T}).$$

References

- [1] S. N. PREOBRAZHENSII, *A consequence of Littlewood's conditional estimates for the Riemann zeta-function*, [arXiv:1104.1358v3](#) [math.NT] (2011).
- [2] Y. MOTOHASHI, *An observation on the zero-free region of the Riemann zeta-function*, *Periodica Mathematica Hungarica* **42** (1–2) (2001), 117–122.
- [3] K. SOUNDARARAJAN, *Moments of the Riemann zeta-function*, [arXiv:math/0612106v2](#) [math.NT] (2008).
- [4] A. E. INGHAM, *The Distribution of Prime Numbers*, Reprint, Stechert-Hafner Service Agency, New York and London, 1964.
- [5] A. SELBERG, *Contributions to the theory of the Riemann zeta-function*, *Arch. Math. Naturvid.* **48** (5) (1946), 89–155.
- [6] E. C. TITCHMARSH, *The Theory of the Riemann Zeta-function*, 2nd ed., revised by D. R. Heath-Brown, Oxford Science Publications, Oxford, 1986.